

FOR ANIMAL TREATMENT ONLY

FLUKECARE® + SE

Combination fluke and roundworm drench for cattle and sheep. For the treatment and control of benzimidazole sensitive mature and immature roundworms, lungworms and early immature (including 2-week-old), mature and immature liver fluke (*Fasciola hepatica*) in cattle and sheep.

ACTIVE CONSTITUENTS

120 g/L triclabendazole, 45.3 g/L oxfendazole and 2.4 g/L sodium selenate.

DIRECTIONS FOR USE

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

DOSAGE AND ADMINISTRATION

Cattle and sheep: 1 ml per 10 kg bodyweight orally. The dose of 1 ml/10 kg delivers 12 mg/kg of triclabendazole, 4.53 mg/kg oxfendazole and 0.1 mg/kg of selenium. Cattle in excess of 650 kg bodyweight to be dosed at 5 ml per 50 kg bodyweight. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (bulls, cows, calves, ewes, wethers, rams, lambs). Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drawing into two or more lines based on bodyweight may be appropriate to avoid excess overdosing. Wash equipment after use. Correct drenching technique should be used.

EFFICACY

For the control of benzimidazole-sensitive mature and immature roundworms, lungworms, and early immature (including 2-week-old), immature and mature liver fluke (*Fasciola hepatica*) in cattle and sheep, listed below:

- Cattle: (*Haemonchus placei*) Barber's pole worm, (*Trichostrongylus axei*) Stomach hair worm, (*Ostertagia ostertagi*) Small brown stomach worm including inhibited (hypobiotic) stages, (*Trichostrongylus spp.*) Black scour worm, (*Cooperia spp.*) Small intestinal worm, (*Nematodirus spp.*) Thin necked intestinal worm, (*Oesophagostomum radiatum*) Nodule worm, (*Dictyoacaulus viviparus*) Lungworm, (*Fasciola hepatica*) Early immature (including 2-week-old fluke), immature and mature liver fluke.
- Sheep: (*Haemonchus contortus*) Barber's pole worm, (*Trichostrongylus axei*) Stomach hair worm, (*Teladorsagia ostertagia spp.*) Small brown stomach worm, (*Trichostrongylus spp.*) Black scour worm, (*Cooperia spp.*) Small intestinal worm, (*Nematodirus spp.*) Thin necked intestinal worm, (*Oesophagostomum columbianum*) Nodule worm, (*Oesophagostomum venulosum*) Large bowel worm, (*Chabertia ovina*) Large mouthed bowel worm, (*Dictyoacaulus filaria*) Large lungworm, (*Fasciola hepatica*) Early immature (including 2-week-old fluke), immature and mature liver fluke.

TREATMENT REGIMEN

Re-dosing intervals vary depending on the age of the animals being treated, degree of parasitism, type of worm and climatic conditions. May be used in pregnant animals at standard dose rates.

GENERAL INSTRUCTIONS

In areas where Black Disease occurs, vaccination is recommended. FLUKECARE® + SE is a free-flowing, ready to use product for the control of acute and chronic parasitic infestation (including two-week-old liver fluke) in cattle and sheep. FLUKECARE® + SE contains triclabendazole and oxfendazole, members of the benzimidazole family of drenches. The dose rate for cattle and sheep is 12 mg/kg of triclabendazole and 4.53 mg/kg of oxfendazole and 0.1 mg/kg of selenium. FLUKECARE® + SE is safe for all classes of animals and has a high safety margin when used as recommended. Resistance may develop to any drench. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It's advisable that a resistance

test be conducted regularly when using any parasite treatment. Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

- MEAT (cattle and sheep): Where the interval between dosing is greater than 28 days: Animals producing meat and oal for human consumption must not be sold for slaughter either during treatment or within 28 days of the last treatment. Where the dosing interval is less than 28 days: Animals producing meat and offal for human consumption must not be sold for slaughter either during treatment or within 56 days of the last treatment.
- MILK (cattle and sheep): Milk intended for sale for human consumption must be discarded during treatment and for 35 days following the last treatment. Not to be used within 28 days of calving or lambing.

REGULATORY INFORMATION

Do not handle until all safety precautions have been read and understood. **WARNING:** Harmful if swallowed. May cause an allergic skin reaction. Suspected of damaging fertility or the unborn child. May cause damage to organs. Toxic to aquatic life with long lasting effects. Harmful to the soil environment. Harmful to terrestrial vertebrates.

HANDLING PRECAUTIONS: Wear protective gloves when handling. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Contaminated clothing should not be allowed out of the work place. Wash contaminated clothing before reuse.

FIRST AID

If medical advice is needed, have product container or label at hand. If swallowed: Call a POISON INFORMATION CENTRE 0800 POISON (0800 764 766) or a doctor if you feel unwell. Rinse mouth. Do not breathe fumes. If on skin: Wash hands thoroughly with plenty of soap and water. If skin irritation or rash occurs get medical advice. If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. If eye irritation persists get medical advice.

DISPOSAL

Triple or pressure rinse container including exterior and cap. Dispose of rinsate and any undiluted chemical according to local government requirements.

STORAGE

Store locked up. Store below 30°C in original container, tightly closed, in a cool dry place. Do not freeze. Do not store for prolonged periods in direct sunlight.

Registered pursuant to the ACVM Act 1997, No. A010203. See www.foodsafety.govt.nz for registration conditions.

Registered to and distributed in New Zealand by:

VIRBAC NEW ZEALAND LIMITED
26-30 Maui Street, Pukete,
Hamilton 3200, New Zealand.
Phone: 0800 VIRBAC (0800 847 222)
Visit us at nz.virbac.com



Ask your **Virbac Area Sales Manager** or **veterinarian** for more information

Shaping the future
of animal health

