

# FlukeCare<sup>®</sup> + Se



A very effective double combination of Triclabendazole and Oxfendazole designed for strategic control of liver fluke and activity against sensitive intestinal parasites.

- Treats all 3 stages of liver fluke with superior control of early immature fluke.
- Patented synergistic control of liver fluke ensuring optimal control.

## DESCRIPTION

Contains 12 mg/kg of Triclabendazole, 45.3 mg/kg of Oxfendazole and 0.1 mg/kg of Selenium. Presented in 5 litre backpacks.

## INDICATIONS

For the treatment and control of benzimidazole sensitive mature and immature roundworms, lungworms and early immature (including 2 week old fluke), mature and immature liver fluke (*Fasciola hepatica*) in cattle and sheep.

## MODE OF ACTION

Triclabendazole and oxfendazole both exhibit enterohepatic recycling. This creates repeated re-exposure of parasitic larval stages in the intestinal mucosa and in the liver, leading to better kill rates. Oxfendazole, together with albendazole and fenbendazole, has a longer half-life than other benzimidazoles. Effective concentrations are maintained for an extended period in the plasma and gut, which increases efficacy against immature and arrested larvae and adult nematodes, including lungworms

## DOSAGE AND ADMINISTRATION

- Cattle and sheep: 1 ml/10 kg bodyweight, orally.

### CATTLE DOSE RATE GUIDE

Bodyweight (kg)	Dose (ml)	Doses per 5 litre
Up to 100	1 ml/10 kg	-
101-150	15	333
151-200	20	250
201-250	25	200
251-300	30	166
301-400	40	125
401-500	50	100
501-600	60	83
601-650	65	76

### SHEEP DOSE RATE GUIDE

Bodyweight (kg)	Dose (ml)	Doses per 5 litre
11-15	1.5	3,333
16-20	2	2,500
21-30	3	1,666
31-40	4	1,250
41-50	5	1,000
51-60	6	833
61-70	7	714
71-75	7.5	666

Cattle in excess of 650 kg bodyweight to be dosed at 5 ml/50 kg bodyweight. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by bodyweight in each group (bulls, cows, calves ewes, wethers, rams, lambs). Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines based in bodyweight maybe appropriate, to avoid excess overdosing. Shake well before use.

## TREATMENT REGIMEN

Re-dosing intervals vary depending on the age of the animals being treated, degree of parasitism, type of worm and climatic conditions. May be used in pregnant animals at standard dose rates.

## WITHHOLDING PERIODS

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

- Milk (cattle and sheep): 35 days
- Meat (where dosing interval is greater than 28 days): 28 days
- Meat (where dosing interval is less than 28 days): 56 days

## REGULATORY INFORMATION

Do not handle until all safety precautions have been read and understood. **WARNING:** Harmful if swallowed. May cause an allergic skin reaction. Suspected of damaging fertility or the unborn child. May cause damage to organs. Toxic to aquatic life with long lasting effects. Harmful to the soil environment. Harmful to terrestrial vertebrates. Wear protective gloves when handling. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Contaminated clothing should not be allowed out of the work place. Wash contaminated clothing before reuse. **FIRST AID:** If medical advice is needed, have product container or label at hand. **IF SWALLOWED:** Call a POISON INFORMATION CENTER 0800 POISON (0800 764 766) or a doctor if you feel unwell. Rinse mouth. Do not breathe fumes. **IF ON SKIN:** Wash hands thoroughly with plenty of soap and water. If skin irritates or rash occurs get medical advice. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses if present and ease to do. Continue rinsing. If eye irritates persists get medical advice.

KEEP OUT OF REACH OF CHILDREN.



## STORAGE

Store locked up. Store below 30°C in original container, tightly closed, in a cool dry place. Do not freeze. Do not store for prolonged periods in direct sunlight.

Registered to and distributed by:

**VIRBAC NEW ZEALAND**

26-30 Maui Street, Pukete, Hamilton.

Visit us at [nz.virbac.com](http://nz.virbac.com)