

Intracillin® 1000 Milking Cow



Intracillin® 1000 Milking Cow is an intramammary mastitis treatment for lactating dairy cows that contains Procaine penicillin G. It's the first choice to treat *Strep. uberis* at calving, and during lactation when a shorter time out of the vat is desired.

- Short 12 hourly treatment intervals and a TAD milk withholding period of only 8 milkings.
- Registered extended therapy claim of 6 tubes given 12 hours apart.

DESCRIPTION

Each single-dose syringe contains 1000 mg of Penicillin. Presented in boxes of 21 syringes.

INDICATION

For the treatment of mastitis infections in lactating cows.

MODE OF ACTION

The high dose of penicillin in this product makes it a very effective treatment for the majority of cases of mastitis in dairy cattle caused by Gram-positive bacteria. Particularly effective against *Strep. uberis*, which is responsible for up to 75% of spring mastitis in New Zealand.

CONTRAINDICATIONS

Not for the treatment of infections caused by organisms that are not susceptible, or have developed resistance, to penicillin. Not to be used in animals with a known history of sensitivity to penicillin-containing products.

DOSAGE & ADMINISTRATION

The entire contents of one syringe should be infused into the teat canal of each infected quarter immediately following milking. Repeat the treatment twice more at 12 hour intervals - three syringes provide a full course of treatment. If required, an extended treatment of 6 syringes (one after each of 6 successive milkings) may be given.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

- Milk: 96 hours / 8 milkings
- Meat: 10 days

REGULATORY INFORMATION

WARNING: May cause eye irritation. Avoid contact with the eyes. May cause sensitisation on prolonged or repeated skin contact, leading to allergic reactions, which may be severe. When handling, wear chemically resistant gloves and avoid skin contact. Penicillins may cause hypersensitivity (allergy) following injection, ingestion or skin contact. Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations. Handle this product with great care to avoid exposure, taking all recommended precautions. **FIRST AID:** If medical advice is needed, have product container or label at hand. **IF SWALLOWED:** Call a POISON INFORMATION CENTRE 0800 POISON (0800 764 766) or a doctor if you feel unwell. Rinse mouth. **IF ON SKIN:** Wash with plenty of soap and water. If skin irritation or rash occurs - get medical advice.

KEEP OUT OF REACH OF CHILDREN.

STORAGE

Store below 25°C. Do not refrigerate.

Registered to and distributed by:

VIRBAC NEW ZEALAND

26-30 Maui Street, Pukete, Hamilton.

Visit us at nz.virbac.com



This product is classified as **green** under Antibiotic Judicious Use Guidelines for the New Zealand Veterinary Profession. Green light antimicrobials are for first line therapy under therapeutic conditions. Visit nzva.org.nz/amr or talk to your Vet to learn more. Please use antibiotics responsibly.

Restricted Veterinary Medicine. Available only under Veterinary Authorisation. Registered Pursuant to the ACVM Act 1997, ACVM No. A7787.

Shaping the future
of animal health

