

KEEP OUT OF REACH OF CHILDREN  
FOR ANIMAL TREATMENT ONLY

# MULTIMIN<sup>®</sup> Copper Free

An injectable supplemental source of zinc, manganese and selenium providing the minerals zinc and manganese in a chelated form for cattle and sheep.

## ACTIVE CONSTITUENTS

Zinc 40 mg/ml (as disodium zinc EDTA), manganese 10 mg/ml (as disodium manganese EDTA), selenium 5 mg/ml (as sodium selenite).

## DIRECTIONS FOR USE

**By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.** For use in cattle and sheep by subcutaneous injection. READ ENTIRE CARTON BEFORE USE.

**Only treat animals in good health. Minimise stress to animals during handling for administration.**

Slight local reaction may occur for about 30 seconds after injection. A slight swelling may be observed at the injection site for a few days after administration. Use standard aseptic procedures during administration of injections. Doses greater than 7 ml should be administered in separate sites. Inject MULTIMIN<sup>®</sup> Copper Free on the opposite side of the neck to any other products or, where this is not possible, at least 10 cm from any other injection sites.

## DOSAGE RECOMMENDATIONS

- Calves (up to 1 year): 1 ml/50 kg
- Cattle (1-2 years): 1 ml/75 kg
- Cattle (over 2 years): 1 ml/100 kg
- Sheep: 0.2 ml/10 kg
- Lambs: 0.2 ml/10 kg
  
- Cows: 3-4 weeks before calving, mating and drying-off.
- Calves: At day 1 and weaning.
- Heifers: Every 3 months, especially 3-4 weeks before mating.
- Rams: 3 times a year.
- Ewes: 4 weeks before tupping, 4 weeks before lambing.
- Lambs: At weaning. As administration of small volumes may lead to dosing errors, it is not recommended to treat lambs under 25kg bodyweight.
- Replacement ewes: Every 3 months - especially 4 weeks before tupping.
- Additional: Every 2 months in wet conditions.

## WITHHOLDING PERIODS

No withholding period is required when used as directed.

## CONTRAINDICATIONS

Do not use at the same time as any other selenised fertiliser, prill, or product without consulting a veterinarian. Do not exceed the stated dose. Do not administer to emaciated animals of body condition score less than 3/10.



## WARNING

Harmful if swallowed. Causes serious eye irritation. May cause damage to organs through prolonged or repeated exposure. Harmful to aquatic life with long lasting effects. IF SWALLOWED: Immediately call a Poison Information centre 0800 POISON (0800 764 766) or a doctor. Rinse mouth. Causes mild skin irritation and serious eye irritation. If skin or eye irritation occurs: Get medical advice. Wear eye protection. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. May cause damage to the liver through prolonged or repeated exposure. Do not breathe spray/mist. Get medical advice if you feel unwell. If medical advice is needed, have product container or label at hand. Avoid release to the environment. Collect spillage. Store locked up.

## STORAGE

Store in original container in a cool, dry place, out of direct sunlight and away from stockfeed or other foodstuffs.

MULTIMIN<sup>®</sup> NZ Patent Number: 524848.

Registered pursuant to the ACVM Act 1997, No. A9882.

See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

Registered to and distributed in New Zealand by:  
**VIRBAC NEW ZEALAND LIMITED**  
26-30 Maui Street, Pukete,  
Hamilton 3200, New Zealand.  
Phone: 0800 VIRBAC (0800 847 222)  
Visit us at [nz.virbac.com](http://nz.virbac.com)

Ask your **Virbac Area Sales Manager** or **veterinarian** for **more information**

Shaping the future  
of animal health

