

# Multimin<sup>®</sup> + Cu



Multimin<sup>®</sup> + Cu is a unique combination of Selenium, Copper, Zinc & Manganese. It is designed to maintain these elements in an animal's optimum range and thereby remove barriers to peak production.

- Boosts innate immunity and response to vaccines.
- Assists with repair and health of reproductive organs.
- Aid in preparing animals for growth.

## DESCRIPTION

An injectable supplemental source of Zinc (40 mg/ml), Manganese (10 mg/ml), Copper (15 mg/ml), and Selenium 5 mg/ml providing the minerals zinc, copper and manganese in a chelated form for cattle and deer. Presented in 200 ml and 500 ml plastic pillow packs.

## INDICATION

To optimise the trace element status of cattle & deer at critical times.

## CONTRAINDICATIONS

Not to be used concurrently with any other form of copper supplementation, or administered at the same time as any other treatment e.g. drenching and vaccination. Do not administer to animals suffering from liver disease, fascioliasis or which have been grazing on plants which may cause liver disease. Only treat animals in good health. Minimise stress to animals during handling for administration. Slight local reaction may occur for about 30 seconds after injection. A slight swelling may be observed at injection site for a few days after administration. Use standard aseptic procedures during administration of injections. Intramuscular injection must be given into the anterior half of the neck. Do not use at the same time as any other selenised fertiliser, prill, or product without consulting a veterinarian. Do not exceed the stated dose.

## DOSAGE & ADMINISTRATION

**By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.**

For use in cattle by subcutaneous or intramuscular injection and in deer by intramuscular injection.

- Calves (up to 1 year): 1 ml/50 kg
- Cattle (1-2 years): 1 ml/75 kg
- Cattle (over 2 years): 1 ml/100 kg
- Deer: 1 ml/100 kg

## TREATMENT REGIMEN

- Cows: 3-4 weeks before calving, breeding and drying off

- Calves: At day 1 and weaning.
- Heifers: Every 3 months, especially 3-4 weeks before mating.
- Additional: Every 2 months in wet conditions.
- Deer: Not recommended more frequently than once every 4 months.

## WITHHOLDING PERIODS

- Milk: Nil
- Meat: Nil

## REGULATORY INFORMATION

**WARNING:** Harmful if swallowed. Wash hands thoroughly after handling. Do not eat, drink or smoke while using this product. **IF SWALLOWED:** Immediately call a Poison Information centre 0800 POISON (0800 764 766) or a doctor. Rinse mouth. Causes mild skin irritation and serious eye irritation. If skin or eye irritation occurs - get medical advice. Wear eye protection. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing. May cause damage to the liver through prolonged or repeated exposure. Do not breathe spray/mist. Get medical advice if you feel unwell. If medical advice is needed, have product container or label at hand. Harmful to aquatic life with long lasting effects. Toxic to terrestrial vertebrates. Avoid release to the environment. Collect spillage. Store locked up.

KEEP OUT OF REACH OF CHILDREN.

## STORAGE

Store in original container in a cool dry place.

Registered to and distributed by:

**VIRBAC NEW ZEALAND**

26-30 Maui Street, Pukete, Hamilton.

Visit us at [nz.virbac.com](http://nz.virbac.com)