

Penclox™ 1200



Penclox™ 1200 is an intramammary mastitis treatment for lactating dairy cows that contains both Procaine penicillin G and cloxacillin. This unique combination makes the product ideal to treat the two most common mastitis pathogens in NZ - *Strep. uberis* & *Staph. aureus*, meaning one product can be used throughout the entire lactation.

- Developed, trialled and made in New Zealand.

DESCRIPTION

Each single-dose syringe contains 1000 mg of Penicillin and 200 mg of Cloxacillin. Presented in boxes of 21 syringes.

INDICATION

For the treatment of mastitis caused by penicillin and/or cloxacillin-sensitive organisms in dairy cows during lactation.

CONTRAINDICATIONS

Not to be used in the treatment of infections caused by organisms known to be resistant to penicillin and cloxacillin. Not to be used in animals with a known history of sensitivity to penicillin or its analogues. Milk from treated cows should not be fed to bobby calves during the treatment or withholding period.

DOSAGE & ADMINISTRATION

The entire contents of one syringe should be infused into the teat canal of each infected quarter immediately following milking. Repeat the treatment twice more at 24 hour intervals - three syringes provide a full course of treatment. If required, an extended treatment of 4, 5, or 6 syringes at 24 hour intervals may be given.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

- Milk (twice-a-day): 108 hours / 9 milkings
- Milk (once-a-day): 120 hours / 5 milkings
- Meat: 10 days

REGULATORY INFORMATION

WARNING: Causes mild skin irritation. May cause an allergic skin reaction. Read label before use. Contaminated work clothing should not be

allowed out of the workplace. Wear protective gloves when handling. Wash contaminated clothing before use. Penicillins may cause hypersensitivity (allergy) following injection, ingestion or skin contact. Do not handle this product if you know you are sensitised, or if you have been advised not to work with such preparations. Handle this product with great care to avoid exposure, taking all recommended precautions. **FIRST AID:** If medical advice is needed, have product container or label at hand. **IF SWALLOWED:** Call a POISON INFORMATION CENTER 0800 POISON (0800 764 766) or a doctor if you feel unwell. Rinse mouth. **IF ON SKIN:** Wash with plenty of soap and water. If skin irritation or rash occurs - get medical advice.

KEEP OUT OF REACH OF CHILDREN.

STORAGE

Store below 25°C. Do not refrigerate.

Registered to and distributed by:

VIRBAC NEW ZEALAND

26-30 Maui Street, Pukete, Hamilton.

Visit us at nz.virbac.com



This product is classified as **orange** under Antibiotic Judicious Use Guidelines for the New Zealand Veterinary Profession. Orange light antimicrobials are restricted to specific indications or used as second line therapy under therapeutic conditions. Visit nzva.org.nz/amr or talk to your Vet to learn more. Please use antibiotics responsibly.

Restricted Veterinary Medicine. Available only under Veterinary Authorisation. Registered Pursuant to the ACVM Act 1997, ACVM No. A10884.

Shaping the future
of animal health

