

VIBROSTREP®

For the treatment of vibriosis (campylobacteriosis), leptospirosis, actinobacillosis (Woody Tongue) and other infections caused by organisms susceptible to streptomycin and dihydrostreptomycin.

ACTIVE CONSTITUENTS

Each ml contains streptomycin (as sulphate) 250 mg and dihydrostreptomycin (as sulphate) 250 mg.

DOSAGE AND ADMINISTRATION

For the treatment of bovine vibriosis:

- Cows: Infuse 2-10 ml into the uterus and inject 3-4 ml per 100 kg bodyweight by subcutaneous or intramuscular injection daily for 3 days.
- Bulls: Infuse and massage 2-10 ml into the preputial cavity and inject 3-4 ml per 100 kg bodyweight by subcutaneous or intramuscular injection daily for 3 days.

For the treatment of vibriosis (campylobacteriosis), leptospirosis, actinobacillosis (Woody Tongue) and other infections caused by organisms susceptible to streptomycin and dihydrostreptomycin:

- Pigs, sheep and cattle: Inject 3-4 ml per 100 kg bodyweight by subcutaneous or intramuscular injection daily for 3 days. In food producing animals, injection must be in the anterior half of the neck.

WITHHOLDING PERIODS

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

- MEAT: Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 30 days of cessation of the last treatment.
- MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than 6 milkings or approximately 72 hrs following the last treatment.

CONTRAINDICATIONS

Not for use in Bobby calves. Not to be used in the treatment of infections caused by organisms known to be resistant to streptomycin or dihydrostreptomycin. Not recommended for use in animals with renal dysfunction or which are already being treated with aminoglycoside (-mycins) antibiotics.

WARNINGS

STERILE INJECTION. KEEP OUT OF REACH OF CHILDREN.

DANGER

Toxic in contact with skin. May damage fertility or the unborn child. Causes damage to hearing. Do not handle until all safety precautions have been read and understood. Wear protective gloves and other personal protective equipment as required. Remove immediately all contaminated clothing. Wash contaminated clothing before reuse. Contaminated work clothing should not be allowed out of the workplace. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. IF ON SKIN: Wash with plenty of soap and water. If exposed or concerned or if a rash occurs: Get medical advice. Call a POISON CENTRE or physician if you feel unwell. Very toxic to aquatic life. Toxic to terrestrial vertebrates. Store locked up. Avoid release to the environment. Collect spillage. Avoid contamination of any water supply with product or empty container. Contains 0.1-1.0 % w/v sodium metabisulfite.



DISPOSAL

Dispose of the empty container in an approved landfill or other approved facility.

STORAGE

Protect from direct sunlight. Store in a cool place.

Registered pursuant to the ACVM Act 1997, No. A3144.

See www.foodsafety.govt.nz for registration conditions.

Registered to and manufactured in New Zealand by:

VIRBAC NEW ZEALAND LIMITED

26-30 Maui Street, Pukete,
Hamilton 3200, New Zealand.
Phone: 0800 VIRBAC (0800 847 222)
Visit us at nz.virbac.com